Depressione PostPartum

*Un'indagine sulle attitudini, le
conoscenze e le pratiche
dei Pediatri di Famiglia
della Provincia di Reggio Emilia*

Annarita Di Buono, Paola Tamburini
(da un progetto di Costantino Panza)

Formazione sulla DPP

Specialità	4	
Corsi Asl	5	
Convegni – Congressi	5	
Monografie	17	
Non ho fatto formazione	40	64%

**La prevalenza della depressione postpartum
materna è:**

2%

5%

8%

13%

La prevalenza della depressione postpartum materna è:

Prevalenza 12.9% primo anno postpartum

Incidenza 14.5% un nuovo episodio nel primo trimestre

International Review of Psychiatry

1996, Vol. 8, No. 1, Pages 37-54

Rates and risk of postpartum depression— a meta-analysis

Michael W. O'hara^{1†} and *Annette M. Swain*¹

The average prevalence rate of non-psychotic postpartum depression based on the results of a large number of studies is **13%**.

Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com

**EUROPEAN
PSYCHIATRY**

European Psychiatry 24 (2009) 327–333

Original article

Prevalence and risk factors for a high level of postnatal
depression symptomatology in Italian women: A sample drawn
from ante-natal classes

Pietro Grussu ^{a,b,*}, Rosa Maria Quatraro ^c

Results. We found that 13% of the women studied showed high postnatal depressive symptomatology

Quante neo-mamme depresse entrano nell'ambulatorio del pediatra in un anno?

3 24

6 27

10

9

14%

Fattori di rischio per una depressione postpartum

Giovane età

Precedenti sintomi depressivi

Immigrata

Assenza del partner

Gravidanza gemellare

Madre > 37 aa

Fecondazione assistita

Postpartum depression

Teri Pearlstein, MD; Margaret Howard, PhD; Amy Salisbury, PhD; Caron Zlotnick, PhD

APRIL 2009 American Journal of Obstetrics & Gynecology

Psychosocial risk factors for PPD include MDD during pregnancy, anxiety during pregnancy, previous nonpuerperal MDD, previous premenstrual dysphoria, stressful life events during pregnancy or the early puerperium, poor social support, marital conflict, low income, immigrant status, and young maternal age.^{14,15} A recent study identified previous depression, current depression and anxiety, and low partner support as key risk factors.¹⁶

Depressione

Marito

Immigrata

Giovane

Fattori di rischio per una depressione postpartum

Giovane età

25

Precedenti sintomi depressivi

58

Immigrata

12

19%

Assenza del partner

43

Gravidanza gemellare

14

Madre > 37 aa

14

Fecondazione assistita

4

Depressione paterna: prevalenza

2%

42

4%

12

10%

2

Paternal depression in the postnatal period and child development: a prospective population study

Paul Ramchandani, Alan Stein, Jonathan Evans, Thomas G O'Connor, and the ALSPAC study team*

Lancet 2005; 365: 2201-05

	% high scores		Crude OR (95% CI)	Adjusted OR (n=5939)*
	Depressed	Not depressed		
Maternal depression (n=9507)				
Emotional	21.9%	12.1%	2.03 (1.71-2.41)	2.03 (1.61-2.56)
Conduct	20.7%	10.7%	2.18 (1.82-2.60)	2.05 (1.60-2.63)
Hyperactivity	12.5%	6.8%	1.97 (1.59-2.45)	1.68 (1.24-2.28)
Prosocial	10.9%	8.2%	1.09 (0.90-1.57)	1.09 (0.79-1.50)
Total	20.7%	7.6%	3.09 (2.6-3.69)	2.76 (2.14-3.55)
Paternal depression (n=7030)				
Emotional	18.5%	12.9%	1.52 (1.10-2.11)	1.27 (0.87-1.87)
Conduct	16.5%	10.6%	1.66 (1.18-2.35)	1.84 (1.27-2.68)
Hyperactivity	11.2%	7.1%	1.65 (1.10-2.47)	1.68 (1.08-2.62)
Prosocial	10.8%	9.1%	1.22 (0.81-1.84)	1.24 (0.78-1.98)
Total	16.9%	8.5%	2.19 (1.55-3.08)	2.09 (1.42-3.08)

16.9%

*Adjusted for effects of social class, degree of maternal education, and depression in other parent.

Table 1: Parental postnatal depressive symptoms and emotional and behavioural problems in children aged 42 months

Individual and Combined Effects of Postpartum Depression in Mothers and Fathers on Parenting Behavior

James F. Paulson, PhD^a, Sarah Dauber, PhD^a, Jenn A. Leiferman, PhD^b

Pediatrics 2006;118;659-668

In this national sample, 14% of mothers and 10% of fathers exhibited levels of depressive symptoms

Nella definizione di depressione postpartum:

Insorge entro la seconda settimana **16**

Insorge entro la sesta settimana **39**

Non si presenta dopo il sesto mese **5**

ICD-10 *International Classification of Diseases*

**F53 – Esordio entro
sei settimane dopo il parto**

DSM – IV

*Diagnostic and Statistical Manual
of Mental Disorders*

**Esordio nelle quattro settimane
dopo un parto**

Investighiamo su una sintomatologia depressiva del genitore ?

Test di “Screening”

Domande di Whooley – *NICE Guideline*

- During the past month have you often been bothered by feeling down, depressed, or hopeless?
 - During the past month have you often been bothered by having little interest or pleasure in doing things?
- If the answer to either of these questions is positive, a third question is recommended by NICE
- Is this something you feel you need or want help with?

Durante gli ultimi mesi, Lei è stata spesso disturbata dal sentirsi giù, dal sentirsi depressa, da una specie di disperazione?

Durante gli ultimi mesi, Lei si è sentita spesso disturbata dal fatto di sentire poco interesse o poco piacere nel fare le cose?

E poi, se la risposta a una di queste domande è positiva:

C'è qualcosa che Lei pensa che possa aiutarla?

Test di “Screening”

Edinburgh Postnatal Depression Scale (EPDS)

SCALA DI DEPRESSIONE PERIPARTUM DI EDIMBURGO
(J. L. Cox, J. K. Olden, R. Sagowsky)

Gentile signora, la invitiamo a scegliere, tra le diverse risposte, senza aiuto e senza discuterne con altre persone, quella che meglio corrisponde al suo stato d'animo negli ultimi sette giorni.

1) Sono stata in grado di ridere e vedere il lato divertente delle cose

2) Mi sono posta in modo positivo verso gli eventi

3) Mi sono sentita colpevole senza motivo quando le cose non andavano bene

4) Sono diventata ansiosa o preoccupata senza motivo

5) Mi sono sentita spaventata o nel panico senza un buon motivo

6) Mi sono sentita sopraffatta dalle cose che accadevano

7) Sono così infelice che ho difficoltà a dormire

8) Mi sono sentita triste o avvilita

9) Sono così infelice che ho pianto

10) Il pensiero di farmi del male mi è venuto in mente

Test di “Screening”

E Lei, signora, come sta?

Qualcuno la aiuta?

Riesce a lasciare il bambino ad altri in qualche momento?

E' un compito del pediatra?

SI **26**

NO **12**

NON SO **24**

36

41%

57%

Choosing a Postpartum Depression Screening Instrument for Your Pediatric Practice

Jacqueline D. Rychnovsky, PhD, RN, CPNP, &
Margaret A. Brady, PhD, RN, CPNP

SUMMARY

The question posed to pediatric outpatient practices caring for new mother/infant dyads is no longer “Should we screen?” but “How should we screen?” new mothers for PPD. PPD screening can be quickly integrated into all well-child visits during the first year of life at minimal cost and administrative burden. Providers must be well connected and familiar with referral sources within their institution or community so that mothers with positive screens receive rapid and effective treatment tailored to their individual needs and symptomatology.

J Pediatr Health Care. (2008). 22, 64-67

Quali ostacoli?

Mancanza di tempo

34

54%

Non sufficiente conoscenza

41

65%

Il genitore non parla

21

Difficoltà di invio

9

Altro

3

Blues materno

Periodo prodromico della DPP **10**

Umore triste dei primi giorni **35**

Sinonimo di DPP **5**

Non risponde

13

Tra i sintomi tipici dello stato depressivo sono presenti

euforia

insonnia

perdita di interesse o piacere nel fare le cose

pensiero di suicidio

allucinazioni

sentimento di colpa o di vergogna

emicrania

Sintomi tipici della DPP

Sentimento di colpa 42

Insonnia 45

**Perdita di interesse e
piacere nel fare 55**

Suicidio 25

ICD-10 *International Classification of Diseases*

Group A symptoms

Depressed mood to a degree that is abnormal for the person, present for most of the day, largely uninfluenced by circumstances

Loss of interest or pleasure in activities that are normally pleasurable

Decreased energy or increased fatiguability

ICD-10 *International Classification of Diseases*

Group B symptoms

Loss of confidence or self esteem

Unreasonable feelings of self reproach or excessive and inappropriate guilt

Recurrent thoughts of death or suicide, or any suicidal behaviour

Complaints or evidence of reduced ability to think or concentrate, such as indecisiveness

Change in psychomotor activity, with agitation or retardation

Sleep disturbance of any type

Change in appetite (decrease or increase) with corresponding weight change

Terapie validate EBM

Psicoterapia cognitiva

24

Terapia farmacologica

24

Nessuna

28

44%

Altro

6

Terapie validate EBM

Psychosocial and psychological interventions for treating postpartum depression (Review)

Dennis C-L, Hodnett E

Authors' conclusions

Although the methodological quality of the majority of trials was, in general, not strong, the meta-analysis results suggest that psychosocial and psychological interventions are an effective treatment option for women suffering from postpartum depression. The long-term effectiveness remains unclear.

United States
National Library
of Medicine

TOXNET

Toxicology Data Network

[TOXNET PDA Access](#)

[SIS Home](#)

[About Us](#)

[Site Map & Search](#)

[Contact Us](#)

[Env. Health & Toxicology](#) [TOXNET](#) [LactMed](#)

Drugs and Lactation Database (LactMed) - A peer-reviewed and fully referenced database of drugs to which breastfeeding mothers may be exposed. Among the data included are maternal and infant levels of drugs, possible effects on breastfed infants and on lactation, and alternate drugs to consider.

Select Database

- [ChemIDplus](#)
- [HSDB](#)
- [TOXLINE](#)
- [CCRIS](#)
- [DART](#)
- [GENETOX](#)
- [IRIS](#)
- [ITER](#)
- **[LactMed](#)**
- [Multi-Database](#)
- [TRI](#)
- [Haz-Map](#)
- [Household Products](#)

Search LactMed

(e.g. Advil, oral contraceptives, Prozac)

For chemicals, add synonyms and CAS numbers to search:
 Yes No

Env. Health & Toxicology

Portal to
environmental
health and
toxicology
resources

[VISIT SITE](#)

Support Pages

- ▶ [LactMed Record Format](#)
- ▶ [Database Creation & Peer Review Process](#)
- ▶ [Help](#)
- ▶ [Fact Sheet](#)
- ▶ [Sample Record](#)
- ▶ [TOXNET FAQ](#)
- ▶ [Glossary](#)
- ▶ [Breastfeeding Links](#)

ADDITIONAL RESOURCE

- [CPDB](#)

[Copyright](#) [Privacy](#) [Accessibility](#)
U.S. National Library of Medicine, 8600 Rockville Pike, Bethesda, MD 20894
National Institutes of Health, Health & Human Services

Last Modified Sun Jul 5 14:05:43 2009

<http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?LACT>

PAROXETINA

Because of the low levels of **paroxetine** in breastmilk, amounts ingested by the infant are small and **paroxetine** has not been detected in the serum of most infants tested. Occasional mild side effects have been reported, especially in the infants of mothers who took **paroxetine** during the third trimester of pregnancy, but the contribution of the drug in breastmilk is not clear. Most authoritative reviewers consider **paroxetine** one of the preferred antidepressants during breastfeeding.[1][2][3][4]

LactMed

SERTRALINA

Because of the low levels of **sertraline** in breastmilk, amounts ingested by the infant are small and is usually not detected in the serum of the infant, although the weakly active metabolite desmethylsertraline is often detectable in low levels in infant serum. Most authoritative reviewers consider **sertraline** one of the preferred antidepressants during breastfeeding.[1][2][3][4]

LactMed

Altri antidepressivi considerati in occasione dell'allattamento al seno

Fluvoxamina, Escitalopram, Fluoxetina, Duloxetina, Citalopram, Maprotilina, Nortryptilina, Phenzelzine, Trazodone, Desipramina, Tranylcypramine, Amitriptylina, Imipramina, Doxepin, Trimipramina, Mirtazapine, Nefazodone, Amoxapina, Velafaxina, Clomipramina, Desvenlafaxina, Olanzapina, Droperidol, Protriptyline,.....

Quale percorso utilizzi per indirizzare la mamma con sintomi depressivi

percorso dedicato aziendale (ASL)	8	
invio al medico di medicina generale	31	49%
invio diretto ad uno psichiatra di fiducia	4	
invio alla NPI	1	
non conosco un percorso definito	21	33%
non attivo alcun percorso		
altro	1	

Quante mamme hai indirizzato negli ultimi dodici mesi verso un percorso dedicato?

Hai mai individuato bambini con patologie o sintomi derivanti da una patologia depressiva del genitore che hanno avuto necessità di intervento medico

si

24

39%

no

37

61%

Quali, tra questi disturbi dell'età pediatrica, si possono associare alla depressione materna

aggressività	3
deficit cognitivo	1
deficit di attenzione	1
tutte le precedenti	53
nessuna delle precedenti	2

88%

The impact of maternal depression and personality disorder on early infant care

Susan Conroy · Maureen N. Marks ·
Robin Schacht · Helen A. Davies · Paul Moran

Neglect

Conclusion Women with depression are less likely to use recommended infant care practices only when they also meet criteria for co-morbid personality disorder. Professionals working with women and babies need to consider the particular difficulties that mothers with both personality disorder and depression may have in providing satisfactory infant care.

“Why didn't somebody warn me?”

The #1 complication of childbirth is depression.

One in eight new moms have postpartum depression.
It is treatable! Call 1-800-944-4PPD.

Extreme exhaustion, appetite and sleep disturbances, mood swings, anxiety, no motivation

Call your health care provider and contact us:
Postpartum Support International
1-800-944-4773
www.postpartum.net